

Picard Groups of the Stable Module Category for Quaternion Groups

Richard Wong
(joint with Jeroen Van de Meer)

2021 Graduate Student Topology and Geometry Conference

Slides can be found at
<http://www.ma.utexas.edu/users/richard.wong/>

Advertisements:

- ▶ electronic Computational Homotopy Theory (eCHT) seminars.
- ▶ Graduates Reminisce Online On Topology (GROOT) Summer Seminar.
- ▶ UT Austin Summer Minicourses - let me know if you're interested in giving a minicourse!

I use the computational methods of homotopy theory to study the **modular representation theory** of finite groups G over a field k of characteristic p , where $p \mid |G|$.

Definition

The **group of endo-trivial modules** is the group

$$T(G) := \{M \in \text{Mod}(kG) \mid \text{End}_k(M) \cong k \oplus P\}$$

where k is the trivial kG -module, and P is a projective kG -module.

We can understand this group as the Picard group of the **stable module category** $\text{StMod}(kG)$:

$$T(G) \cong \text{Pic}(\text{StMod}(kG))$$

Theorem (van de Meer-W., cf Carlson-Thévenaz)

Let ω denote a cube root of unity.

$$\text{Pic}(\text{StMod}(kQ_8)) \cong \begin{cases} \mathbb{Z}/4 & \text{if } \omega \notin k \\ \mathbb{Z}/4 \oplus \mathbb{Z}/2 & \text{if } \omega \in k \end{cases}$$

Theorem (van de Meer-W., cf Carlson-Thévenaz)

Let $n \geq 4$.

$$\text{Pic}(\text{StMod}(kQ_{2^n})) \cong \mathbb{Z}/4 \oplus \mathbb{Z}/2$$

Definition

The **Picard group** of a symmetric monoidal category $(\mathcal{C}, \otimes, 1)$, denoted $\text{Pic}(\mathcal{C})$, is the set of isomorphism classes of invertible objects X , with

$$\begin{aligned}[X] \cdot [Y] &= [X \otimes Y] \\ [X]^{-1} &= [\text{Hom}_{\mathcal{C}}(X, 1)]\end{aligned}$$

Example (Hopkins-Mahowald-Sadofsky)

For $(\text{Sp}, \wedge, \mathbb{S}, \Sigma)$ the stable symmetric monoidal category of spectra,

$$\text{Pic}(\text{Sp}) \cong \mathbb{Z}$$

Given a symmetric monoidal ∞ -category \mathcal{C} , one can do better than the Picard group:

Definition

The **Picard space** $\mathrm{Pic}(\mathcal{C})$ is the ∞ -groupoid of invertible objects in \mathcal{C} and isomorphisms between them.

This is a group-like E_∞ -space, and so we equivalently obtain the connective **Picard spectrum** $\mathrm{pic}(\mathcal{C})$.

Proposition (Mathew-Stojanoska)

The functor $\mathrm{pic} : \mathrm{Cat}^\otimes \rightarrow \mathrm{Sp}_{\geq 0}$ commutes with limits and filtered colimits.

Example

Let R be an E_∞ -ring spectrum. Then $\mathrm{Mod}(R)$ is a stable symmetric monoidal ∞ -category.

The homotopy groups of $\mathrm{pic}(R) := \mathrm{pic}(\mathrm{Mod}(R))$ are given by:

$$\pi_*(\mathrm{pic}(R)) \cong \begin{cases} \mathrm{Pic}(R) & * = 0 \\ (\pi_0(R))^\times & * = 1 \\ \pi_{*-1}(\mathfrak{gl}_1(R)) \cong \pi_{*-1}(R) & * \geq 2 \end{cases}$$

Definition

The **stable module category** $\text{StMod}(kG)$ has objects kG -modules, and has morphisms

$$\underline{\text{Hom}}_{kG}(M, N) = \text{Hom}_{kG}(M, N) / \text{PHom}_{kG}(M, N)$$

where $\text{PHom}_{kG}(M, N)$ is the linear subspace of maps that factor through a projective module.

Proposition

$\text{StMod}(kG)$ is a stable symmetric monoidal ∞ -category.

From now on, we restrict our attention to the case that G is a finite p -group, so that the following theorem holds:

Theorem (Keller, Mathew, Schwede-Shipley)

There is an equivalence of symmetric monoidal ∞ -categories

$$\mathrm{StMod}(kG) \simeq \mathrm{Mod}(k^{tG})$$

Where k^{tG} is an E_∞ ring spectrum called the G -Tate construction.

Definition

The **G -Tate construction** is the cofiber of the norm map:

$$k_{hG} \xrightarrow{N_G} k^{hG} \rightarrow k^{tG}$$

Theorem (Mathew-Stojanoska)

If $f : R \rightarrow S$ is a **faithful G -Galois extension** of E_∞ ring spectra, then we have an equivalence of ∞ -categories

$$\mathrm{Mod}(R) \cong \mathrm{Mod}(S)^{hG}$$

Corollary

We have the **homotopy fixed point spectral sequence**, which has input the G action on $\pi_*(\mathrm{pic}(S))$:

$$H^s(G; \pi_t(\mathrm{pic}(S))) \Rightarrow \pi_{t-s}(\mathrm{pic}(S)^{hG})$$

whose abutment for $t = s$ is $\mathrm{Pic}(R)$.

Definition

A map $f : R \rightarrow S$ of E_∞ -ring spectra is a **G -Galois extension** if the maps

(i) $i : R \rightarrow S^{hG}$

(ii) $h : S \otimes_R S \rightarrow F(G_+, S)$

are weak equivalences.

Definition

A G -Galois extension of E_∞ -ring spectra $f : R \rightarrow S$ is said to be **faithful** if the following property holds:

If M is an R -module such that $S \otimes_R M$ is contractible, then M is contractible.

Proposition (Rognes)

A G -Galois extension of E_∞ -ring spectra $f : R \rightarrow S$ is faithful if and only if the Tate construction S^{tG} is contractible.

Proposition (van de Meer-W.)

For Q a quaternion group with center $H \cong \mathbb{Z}/2$,

$$k^{hQ} \rightarrow k^{h\mathbb{Z}/2} \text{ and } k^{tQ} \rightarrow k^{t\mathbb{Z}/2}$$

are faithful Q/H -Galois extensions of ring spectra.

Lemma

$$\pi_*(k^{h\mathbb{Z}/2}) \cong k[t^{-1}] \quad \pi_*(k^{t\mathbb{Z}/2}) \cong k[t^{\pm 1}]$$

Lemma

Note that $Q_8/H \cong (\mathbb{Z}/2)^2$.


$$H^*((\mathbb{Z}/2)^2; k) \cong k[x_1, x_2] \text{ with } |x_i| = 1.$$

Note that $Q_{2^n}/H \cong D_{2^{n-1}}$.

$$H^*(D_{2^{n-1}}; k) \cong k[x_1, u, z]/(ux_1 + x_1^2 = 0)$$

with $|x_i| = |u| = 1, |z| = 2$. Moreover, $\text{Sq}^1(z) = uz$.

$$E_2^{s,t} = H^s(Q/H; \pi_t(k^{h\mathbb{Z}/2})) \Rightarrow \pi_{t-s}(k^{hQ})$$



The Adams-graded E_2 page. $\circ = k$. Not all differentials are drawn.

Proposition

For $G = Q_8$, we have differentials


$$d_2(t) = x_1^2 + x_1x_2 + x_2^2 \quad \text{and} \quad d_3(t^2) = x_1^2x_2 + x_1x_2^2$$

Proposition

For $G = Q_{2^n}$, we have differentials


$$d_2(t) = u^2 + z \quad \text{and} \quad d_3(t^2) = uz$$

$$E_2^{s,t} = H^s(Q/H; \pi_t(k^{t\mathbb{Z}/2})) \Rightarrow \pi_{t-s}(k^{tQ})$$




The Adams-graded E_2 page. $\circ = k$. Not all differentials are drawn.

$$E_2^{s,t} = H^s(Q/H; \pi_t(k^{t\mathbb{Z}/2})) \Rightarrow \pi_{t-s}(k^{tQ})$$




The Adams-graded E_3 page. $\circ = k$. Not all differentials are drawn

$$E_2^{s,t} = H^s(Q/H; \pi_t(k^{t\mathbb{Z}/2})) \Rightarrow \pi_{t-s}(k^{tQ})$$




The Adams-graded $E_4 = E_\infty$ page. $\circ = k$.

$$E_2^{s,t} = \widehat{H}^s(Q/H; \pi_t(k^{t\mathbb{Z}/2})) \Rightarrow \pi_{t-s}((k^{t\mathbb{Z}/2})^{tQ/H})$$




The Adams graded E_2 page of the Tate spectral sequence. $\circ = k$. Not all differentials are drawn.

$$E_2^{s,t} = \widehat{H}^s(Q/H; \pi_t(k^{t\mathbb{Z}/2})) \Rightarrow \pi_{t-s}((k^{t\mathbb{Z}/2})^{tQ/H})$$



The Adams graded E_4 page of the Tate spectral sequence. $\circ = k$.

$$E_2^{s,t} = \widehat{H}^s(Q/H; \pi_t(k^{t\mathbb{Z}/2})) \Rightarrow \pi_{t-s}((k^{t\mathbb{Z}/2})^t Q/H)$$



The Adams graded E_4 page of the Tate spectral sequence. $\circ = k$.

Proposition (van de Meer-W.)

For Q a quaternion group with center $H \cong \mathbb{Z}/2$,

$$k^{hQ} \rightarrow k^{h\mathbb{Z}/2} \text{ and } k^{tQ} \rightarrow k^{t\mathbb{Z}/2}$$

are faithful Q/H -Galois extensions of ring spectra.

Therefore, we can compute

$$T(Q) \cong \mathrm{Pic}(\mathrm{StMod}(kQ)) \cong \mathrm{Pic}(k^{tQ})$$

Corollary

We have the HFPSS


$$H^s(Q/H; \pi_t(\mathrm{pic}(k^{t\mathbb{Z}/2}))) \Rightarrow \pi_{t-s}(\mathrm{pic}(k^{t\mathbb{Z}/2})^{hQ/H})$$

whose abutment for $t = s$ is $\mathrm{Pic}(\mathrm{StMod}(kQ))$.

Proposition

The homotopy groups of $\mathrm{pic}(k^{t\mathbb{Z}/2})$ are given by:

$$\pi_*(\mathrm{pic}(k^{t\mathbb{Z}/2})) \cong \begin{cases} \mathrm{Pic}(k^{t\mathbb{Z}/2}) \cong 1 & * = 0 \\ k^\times & * = 1 \\ \pi_{*-1}(k^{t\mathbb{Z}/2}) \cong k & * \geq 2 \end{cases}$$



The Adams graded E_2 page of the HFPSS computing $\pi_*(\text{pic}(k^{t\mathbb{Z}/2})^{hQ/H})$. Not all differentials are drawn. $\circ = k$, $\blacklozenge = k^\times$.

By the construction of $\mathrm{pic}(R)$, we have an identification of differentials $d_r^{s,t}(\mathrm{pic}S) \cong d_r^{s,t-1}(S)$ for $t - s > 0$ and $s > 0$.

Theorem (Mathew-Stojanoska)


Let $R \rightarrow S$ be a G -Galois extension of E_∞ ring spectra. Then we further have an identification of differentials for $2 \leq r \leq t - 1$, which yields an isomorphism

$$f : E_t^{t,t-1}(S) \xrightarrow{\cong} E_t^{t,t}(\mathrm{pic}(S))$$


Moreover, there is a formula for the first differential outside of this range:

$$d_t^{t,t}(f(x)) = f(d_t^{t,t-1}(x) + x^2), \quad x \in E_t^{t,t-1}(S)$$


$$E_2^{s,t} = H^s(Q/H; \pi_t(k^{t\mathbb{Z}/2})) \Rightarrow \pi_{t-s}(k^{tQ})$$



The Adams-graded E_2 page. $\circ = k$. Not all differentials are drawn.



The Adams graded E_2 page of the HFPSS computing $\pi_*(\text{pic}(k^{t\mathbb{Z}/2})^{hQ/H})$. Not all differentials are drawn. $\circ = k$, $\blacklozenge = k^\times$.




The Adams graded E_3 page of the HFPSS computing $\pi_*(\text{pic}(k^{tQ_8}))$. Not all differentials are drawn. $\circ = k$, $\blacklozenge = k^\times$, $\bullet = \mathbb{Z}/2$.

For $G = Q_8$, note that $E_3^{3,3} \cong k^2$, generated by $t^{-2}x_1x_2^2$ and $t^{-2}x_1^2x_2$. Applying the formula for the differential for $\alpha, \beta \in k$, noting that $x_1^3x_2^3 = x_1^4x_2^2 + x_1^2x_2^4$, we have

$$\begin{aligned} d_3(f(\alpha t^{-2}x_1^2x_2)) &= f(\alpha t^{-4}(x_1^4x_2^2 + x_1^3x_2^3)) + f(\alpha^2 t^{-4}x_1^4x_2^2) \\ &= f(\alpha t^{-4}(x_1^4x_2^2 + (x_1^4x_2^2 + x_1^2x_2^4))) + f(\alpha^2 t^{-4}x_1^4x_2^2) \\ &= f(\alpha^2 t^{-4}x_1^4x_2^2) + f(\alpha t^{-4}x_1^2x_2^4) \end{aligned}$$

$$\begin{aligned} d_3(f(\beta t^{-2}x_1x_2^2)) &= f(\beta t^{-4}(x_1^3x_2^3 + x_1^2x_2^4)) + f(\beta^2 t^{-4}x_1^2x_2^4) \\ &= f(\beta t^{-4}((x_1^4x_2^2 + x_1^2x_2^4) + x_1^2x_2^4)) + f(\beta^2 t^{-4}x_1^2x_2^4) \\ &= f(\beta^2 t^{-4}x_1^2x_2^4) + f(\beta t^{-4}x_1^4x_2^2) \end{aligned}$$

For an element to be in the kernel, we then must simultaneously have the expressions $\alpha + \beta^2 = 0$ and $\beta + \alpha^2 = 0$.



The Adams graded E_4 page of the HFPSS computing $\text{pic}((k)^{t_{Q_8}})$, where k has a cube root of unity. $\circ = k$, $\bullet = \mathbb{Z}/2$, $\blacklozenge = k^\times$.

For $G = Q_{2^n}$, note that $E_3^{3,3} \cong k^2$, generated by $t^{-2}uz$ and $t^{-2}x_1z$. Applying the formula for the differential for $\alpha, \beta \in k$, noting that $ux_1 = x_1^2$ in the E_3 page, we have

$$\begin{aligned} d_3(f(\alpha t^{-2}uz)) &= f(\alpha u^2 z^2 t^{-4}) + f(\alpha^2 u^2 z^2 t^{-4}) \\ &= f((\alpha + \alpha^2)u^2 z^2 t^{-4}) \end{aligned}$$

$$\begin{aligned} d_3(f(\beta t^{-2}x_1z)) &= f(\beta(ux_1z^2)t^{-4}) + f(\beta^2 x_1 z^2 t^{-4}) \\ &= f((\beta + \beta^2)x_1 z^2 t^{-4}) \end{aligned}$$

For an element to be in the kernel, we must simultaneously have the expressions $\alpha + \alpha^2 = 0$ and $\beta + \beta^2 = 0$.

Theorem (van de Meer-W.)

Let ω denote a cube root of unity.

$$\text{Pic}(\text{StMod}(kQ_8)) \cong \begin{cases} \mathbb{Z}/4 & \text{if } \omega \notin k \\ \mathbb{Z}/4 \oplus \mathbb{Z}/2 & \text{if } \omega \in k \end{cases}$$

Theorem (van de Meer-W.)




Let $n \geq 4$.

$$\text{Pic}(\text{StMod}(kQ_{2^n})) \cong \mathbb{Z}/4 \oplus \mathbb{Z}/2$$

Future Directions

- ▶ **Generalizations** - Compute $\text{Pic}(\text{StMod}(kG))$ for G dihedral and semi-dihedral, or for extraspecial and almost-extraspecial p -groups.
- ▶ **Tensor-Triangulated Geometry** - Compute $\text{Pic}(\Gamma_p(\text{StMod}(kG)))$, where $\Gamma_p(\text{StMod}(kG))$ denotes a thick or localizing tensor-ideal subcategory of $\text{StMod}(kG)$.
- ▶ Categorify the Dade group of endo-permutation modules.
- ▶ Further HFPSS or Tate spectral sequence calculations.

References

-  Carlson, Jon F., and Thévenaz, Jacques.
Torsion Endo-Trivial Modules.
Algebras and Representation Theory 3 (4): 303–35, 2000.
-  Mathew, Akhil, and Stojanoska, Vesna.
The Picard group of topological modular forms via descent theory.
Geom. Topol. 20 (6): 3133–3217, 2016.
-  Mathew, Akhil.
The Galois group of a stable homotopy theory.
Advances in Mathematics 291: 403–541, 2016.